

Conception and development of Sexual Health indicators to advance the quality of spinal cord injury rehabilitation: SCI-High Project

Stacy Elliott, Gaya Jeyathevan, Shea Hocaloski, Colleen O'Connell, Sivakumar Gulasingam, Sandra Mills, Farnoosh Farahani, Anita Kaiser, S. Mohammad Alavinia, Maryam Omidvar & B. Catharine Craven

To cite this article: Stacy Elliott, Gaya Jeyathevan, Shea Hocaloski, Colleen O'Connell, Sivakumar Gulasingam, Sandra Mills, Farnoosh Farahani, Anita Kaiser, S. Mohammad Alavinia, Maryam Omidvar & B. Catharine Craven (2019) Conception and development of Sexual Health indicators to advance the quality of spinal cord injury rehabilitation: SCI-High Project, *The Journal of Spinal Cord Medicine*, 42:sup1, 68-84, DOI: [10.1080/10790268.2019.1612642](https://doi.org/10.1080/10790268.2019.1612642)

To link to this article: <https://doi.org/10.1080/10790268.2019.1612642>



© 2019 The Author(s). Published with license by Taylor & Francis Group, LLC



Published online: 01 Oct 2019.



[Submit your article to this journal](#)



Article views: 460



[View related articles](#)



[View Crossmark data](#)



Citing articles: 1 [View citing articles](#)

Research Article

Conception and development of Sexual Health indicators to advance the quality of spinal cord injury rehabilitation: SCI-High Project

Stacy Elliott^{1,2,3}, Gaya Jeyathevan ⁴, Shea Hocaloski¹, Colleen O'Connell⁵, Sivakumar Gulasingam^{6,7}, Sandra Mills⁷, Farnoosh Farahani ⁴, Anita Kaiser ^{4,8}, S. Mohammad Alavinia ^{4,6}, Maryam Omidvar⁴, B. Catharine Craven ^{4,6,7}

¹Sexual Health Rehabilitation Service, GF Strong Rehabilitation Center & Blusson Spinal Cord Centre, Vancouver, British Columbia, Canada, ²Faculty of Medicine, ICORD, University of British Columbia, Vancouver, British Columbia, Canada, ³BC Center for Sexual Medicine, Vancouver Hospital, Vancouver, British Columbia, Canada, ⁴KITE, Toronto Rehab – University Health Network, Toronto, Ontario, Canada, ⁵Stan Cassidy Centre for Rehabilitation, Fredericton, New Brunswick, Canada, ⁶Department of Medicine, Division of Physical Medicine and Rehabilitation, University of Toronto, Toronto, Ontario, Canada, ⁷Brain and Spinal Cord Rehabilitation Program, Toronto Rehabilitation Institute – University Health Network, Toronto, Ontario, Canada, ⁸Canadian Spinal Research Organization, Toronto, Ontario, Canada

Context: Although spinal cord injury or disease (SCI/D) results in complex biological and psychosocial impairments that adversely impact an individual's overall quality of sexual life, sexual health is poorly integrated into the current rehabilitation processes. Therefore, it is vital to promote sexual health as a rehabilitation priority. Herein, we describe the selection of Sexual Health structure, process and outcome indicators for adults with SCI/D in the first 18 months after rehabilitation admission.

Methods: Experts in sexual health and the SCI-High team identified key factors that influence the sexual health outcomes of rehabilitation interventions to inform Driver diagram development. This diagram informed the selection and development of indicators to promote a permissive environment for discussion of sexual health issues among regulated health care professionals (HCPs). A review of literature and psychometric properties of measurement tools facilitated final indicators selection.

Results: The *structure indicator* is the proportion of rehabilitation HCPs who have completed annual preliminary sexual health training. The *process indicator* is the proportion of SCI/D inpatients that have a documented introduction to available local sexual health resources. The *outcome indicator* is a sexual health patient questionnaire used to assess sexual health patient outcomes and sexual health information/educational needs. Rapid-cycle piloting verified that the indicator tools developed are feasible for implementation.

Conclusion: Successful implementation of the Sexual Health structure, process and outcome indicators will promote a permissive environment to enable open discussion, and lead to provision of equitable and optimal care related to sexual health following SCI/D. This will ultimately advance sexual health rehabilitation across the nation.

Keywords: Spinal cord injuries, Sexual health, Healthcare quality indicators, Rehabilitation

Correspondence to: B. Catharine Craven, Neural Engineering & Therapeutic Team, KITE, Toronto Rehabilitation Institute – University Health Network, 520 Sutherland Drive, Toronto, ON M4G 3V9, Canada; Ph: (416) 597-3422 (ext: 6122). Email: cathy.craven@uhn.ca

Introduction

Sexuality is, and always has been, a major priority for human beings. Beyond the basic necessity for reproduction, sexual activity is a form of social engagement, and a way to connect with others in a mentally, spiritually,

and physically intimate and meaningful way.^{1,2} Although these fundamental motives do not change following spinal cord injury or disease (SCI/D), significant physiologic changes occur with respect to sensation, voluntary motor and autonomic function altering sexual arousal, orgasmic potential, ejaculation in men, and sexual positioning for both sexes.³ Furthermore, the secondary health consequences of spinal cord impairment, including bladder and bowel continence and related issues, autonomic dysreflexia, nociceptive and neuropathic pain, depression, and pharmacologic management of such conditions, also affect sexual self-esteem, sexual expression, and sexual satisfaction. Despite these aforementioned impairments, sexual interest typically continues to be a priority for many. Readiness and courage are therefore required on the part of the individual with SCI/D and their current or future partner(s) to learn how to adapt to the functional changes in the body to experience a rewarding and satisfying sexual life after SCI/D.

For the purposes of this manuscript, *Sexual Self-Esteem* refers to a positive regard for, and confidence in, the capacity to experience one's sexuality in a satisfying and enjoyable way; *Sexual Expression* refers to the way in which we present and communicate as a sexual being to the world, through feelings and behaviors around sexual acts, sexual orientation and gender identity; and, *Sexual Satisfaction* refers to the expression of positive sexual outcomes and relationship intimacy following sexual activity. *Sexual Activity* refers to sexual acts with self or partner(s), which may or may not include penetrative activity. *Sexual Life* refers to the physical, mental, emotional and social aspects of sexual activity, expression and relationships. *Sexual Enjoyment* refers to the pleasure derived from sexual activity with self or partner(s). *Sexual Interest* refers to sexual urge and/or motivation to seek out and/or engage in sexual activity.

It is well established that improving sexual activity and sexual self-esteem improves quality of life after SCI/D.^{4,5} Data from a large Scandinavian survey of 545 women with SCI concluded that 80% of women who continued to be sexually active after injury were able to overcome the physical restrictions and mental obstacles related to their injury, and lead active and positive sexual lives with a partner.⁶ In men, following traumatic SCI, erectile function, ejaculation and orgasm are most severely affected, while sexual desire remains high.⁷ Personal interviews with 33 heterosexual men with chronic SCI demonstrated that a willingness to adapt their sexual behavior, coupled with the reward of physical pleasure experienced during intercourse, can result in a positive and satisfying sex life.⁸ High-

quality SCI rehabilitation training programs for patients should address expectations for sexual activity and sexual satisfaction within their core curriculum.

A common, albeit outdated, myth is that after SCI/D, one ceases to be capable of being sexually active or loses interest in sexual activity, leading to the notion that sex should not be a rehabilitation priority. While other "more comfortable" secondary health conditions, and their related management concerns dominate rehabilitation programs and rehabilitation research, sexual and fertility rehabilitation is, and *always* has been, a major priority for persons with SCI/D. It is imperative that such 'mismatch' of priorities are more thoroughly addressed by health care professionals (HCPs) in rehabilitation settings. Until recently, sexual and fertility rehabilitation has not been a major priority for HCPs involved in the overall physical, medical and psychological rehabilitation following SCI/D. There are many contributing factors to this traditional neglect, including the personal nature and sensitivity of the subject, the discomfort experienced by HCPs when addressing the topic, and a lack of knowledge or confidence regarding the assessment and management of sexual health among HCPs. However, with the advent of Viagra[®] (sildenafil) for erectile dysfunction in the late 1990s, the subject of sex has been broached more openly in various media outlets, and expectations of patients with sexual health concerns have risen. Unfortunately, HCPs involved in SCI/D rehabilitation as a whole have not responded to this need with proportional effort in either clinical or research initiatives⁹ resulting in a major gap in care.

Given that sexual health is biopsychosocial in nature, there are numerous domains in which sexual health issues will arise. This underscores the need for various disciplines (e.g. doctors, nurses, physical and occupational therapists, social workers, psychologists, recreational therapists, etc.) to be informed and competent in providing fundamental sexual and rehabilitation information.³ As a result of substantial variation in clinical services available, coupled with a paucity of formalized sexual rehabilitation services or prioritization of sexual health goals, concerted efforts are required, nationally, to address the current gaps in care and advance sexual health rehabilitation with the goal of improving standards of SCI/D rehabilitation in Canada by 2020.

Indicators can measure the structure, process and outcome of health care services which can evidently facilitate the sustainability of a high-quality health care system.¹⁰ *Structure indicators* are defined by the properties of the setting in which the health care services are delivered¹¹ while *process indicators* describe the

specific activities involved in providing and receiving of care.¹² Finally, *outcome indicators* describe the effects of health care provided to a given individual or population (e.g. mortality, health-related quality of life, or patient satisfaction, etc.).¹² In combination, these indicators serve as a foundation for the development of an evaluation framework that can be widely implemented across various hospital settings as a benchmark for optimizing the overall quality of health care. Indicator data is commonly used to identify gaps and trends in care, inform priority setting and policy formulation, and monitor rehabilitation programs. Furthermore, indicators can facilitate comparisons across and between different health care settings and ensure continuous quality improvement (i.e. benchmarking) while building transparency in health care.^{13,14}

The SCI Rehabilitation Care High Performance Indicators (or “SCI-High”) Project endeavors to advance SCI rehabilitation care by 2020 through consensus derived development, selection, implementation and evaluation of indicators of quality care for 11 Domains of rehabilitation care prioritized by clinicians, researchers and individuals living with chronic SCI/D. This manuscript describes the process involved in the selection, development and implementation of structure, process and outcome indicators for the Sexual Health Domain for adults with SCI/D from rehabilitation admission to 18 months post-admission.

Methods

A detailed description of the overall SCI-High Project methods and process for identifying Sexual Health as a priority Domain for SCI rehabilitation care are described in related manuscripts in this issue.^{15,16} In addition to the investigative team (www.sci-high.ca), an external advisory committee and national data strategy committee supported the global Project goals and provided oversight regarding the context for implementing all of the planned indicators.

The approach to developing the Sexual Health Domain’s structure, process and outcome indicators followed a modified, but substantially similar, approach to that described by Mainz,¹¹ which included planning and development phases: (a) formation and organization of national and local Working Groups;⁹ (b) defining and refining the key Domain and specific target construct; (c) providing an overview/summary of existing evidence and practice; (d) developing and interpreting a Driver diagram; (e) selecting indicators; and (f) pilot testing and refinement of the Domain-specific structure, process and outcome indicators. Throughout these processes, a facilitated discussion occurred amongst the

Domain-specific Working Group and the SCI-High Project Team to utilize relevant expertise on the topic, while ensuring that the broader goals of the SCI-High Project were aligned across the other 10 Domain Working Groups (as appropriate).

Sexual Health Working Group

Experts in sexual health and relevant stakeholders were invited to participate in the SCI-High Project as members of the Domain-specific Working Group based on their practical or empirical knowledge of SCI/D rehabilitation, sexual health, health service delivery and patient education. The group included sexual health/medicine practitioners, patient and family educators, physiatrists, nurses, postdoctoral fellows and a stakeholder with lived experience. The Working Group met nine times via web-enabled conference calls over an 8-month period, totaling 9 h of discussion related to the development of the indicators, and an additional 3 h thereafter, to refine the indicators and discuss manuscript preparation. Outside of the formal meetings, individual members of the Working Group completed an additional review of the prepared materials, shared resources and/or practice standards with one another, and/or conducted independent evaluations of the proposed indicators.

Driver diagram and construct definition

The selection of Sexual Health as a Domain of interest for indicator development emerged from a consensus-building activity intended to select the broader set of Domains being pursued within the overarching SCI-High Project.⁹ This process involved a systematic search to collect information about SCI/D rehabilitation care related to sexual health, identification of factors that influence the outcome of rehabilitation interventions, and a scoping synthesis of the data acquired. MEDLINE, EMBASE and CINAHL databases were searched using the terms “sexual health”, “spinal cord injury”, or both. The group initially developed four Driver diagrams that were sex and gender specific and related to sexual health. Following several cycles of Driver diagram review and refinement, all four were amalgamated into a single driver diagram (Figure 1). A Driver diagram is a visual display of a high-level quality improvement goal, and a set of underpinning factors/goals. The tool helped to organize change concepts as the Working Group discerned “what changes can we make, that will result in goal attainment”. The branches in red within the final Driver diagram represent the main areas that were the focus for the development of indicators based on experts’ opinions.




Figure 1 Sexual Health Driver diagram. The impairment branch is common to the 11 SCI-High Project Domains. UEMS: upper-extremity motor score; LEMS: lower-extremity motor score; NLI: neurological level of injury; AIS: ASIA impairment scale; HR: heart rate; BP: blood pressure; IUD ± H: intrauterine devices ± hormone therapy; PVS: penile vibratory stimulation; MESA: microsurgical epididymal sperm aspiration; TESA: testicular sperm aspiration; PESA: percutaneous epididymal sperm aspiration; ICSI: intracytoplasmic sperm injection; IVF: *in vitro* fertilization; IUI: intrauterine insemination; IVI: intravaginal insemination; PDE5I: phosphodiesterase Type 5 inhibitors; ICI: intracervical insemination; CBT: cognitive behavioral therapy; LESWT: low energy shock wave therapy; EROS: Environmental Rewards Observation Scale; TENS: transcutaneous electrical nerve stimulation; AD: autonomic dysreflexia. *Oral contraception, patches/rings, and provera. **Opioids, antidepressants, antispastic, antihypertensive, and hormones.

Following review of the systematic searches, discussions, and multiple refinements of the Driver diagram, the group agreed that creating a permissive environment among regulated HCPs in relation to sexual health was the driver most likely to advance SCI/D rehabilitation care in the near term. Based on this discussion, and reflection on current terminology related to expression of gender and sexuality, the following construct definition was created: *The World Health Organization (WHO) defines sexual health as a state of physical, emotional, mental and social well-being in relation to sexuality; it is not merely the absence of disease, dysfunction or infirmity. Sexual health requires a positive and respectful approach to sexuality and sexual relationships, as well as the possibility of having pleasurable and safe sexual experiences, free of coercion, discrimination and violence. For sexual health to be attained and maintained, the sexual rights of all persons must be respected, protected and fulfilled.*¹² In this context, sexuality encompasses: sexual activity, gender identity, gender roles, sexual orientation, eroticism, pleasure, intimacy, contraception and reproduction. Sexual health rehabilitation requires a positive and respectful approach to sexuality, self-esteem, sexual relationships, and reproductive wishes, as well as the potential to have consensual, pleasurable and safe sexual experiences.

Selection of indicators

The Working Group asked to develop/select at least one indicator each for structure, process and outcome in relation to the Sexual Health Domain. The Project Leaders stipulated that the indicators should be relevant, concise and feasible to implement (10 min or less), and aligned across the structure, process and outcome to achieve a single substantive advance in SCI/D rehabilitation care. The indicators could be

measured using established or new measurement tools (i.e. questionnaires, data collection sheets, laboratory exams, and medical record data), depending on the requirements and feasibility of a given indicator.

Indicator piloting

In order to create a permissive environment, enabling open discussions and to promote sexual health (including post SCI/D sexual activity, self-esteem, expression and satisfaction), the feasibility of the structure and outcome indicators for the Sexual Health Domain were pilot tested. The indicators were reviewed and refined through multiple quality improvement Plan-Do-Study-Act (PDSA)¹³ cycles for quick qualitative evaluations, feedback and refinement.

Pilot study methods – structure indicator

First, a pre and post-self-assessment tool was developed and implemented at a tertiary SCI rehabilitation hospital to assess the knowledge of HCPs involved in SCI/D rehabilitation and to inform sexual health training and education. HCPs were asked to participate in a day-long sexual health workshop and were provided with a sexual health education video comprised of several modules merged into one YouTube online resource. The video modules were developed by the Spinal Cord Injury Research Evidence (SCIRE) project team and Facing Disability, and focused on approaching/initiating patient discussion on sexual health and directing them to appropriate sexual health resources/services (please see Table 1). Permission to use the video modules in this manner was provided by the SCIRE project team. Prior to and following the video modules and workshop provision, a self-assessment tool was administered to HCPs to in order to assess their acquisition of sexual health knowledge, such as active learning and knowledge retention (structure

Table 1 Sexual health video modules for the structure indicator.

Video module	Title & publisher	YouTube link	Duration
1.	Sexual Health and Spinal Cord Injury: An Introduction By: SCIREWebVideo	https://www.youtube.com/watch?v=H1waqXyrRQI	3:16
2.	3 Sexual Medicine Principles and Spinal Cord Injury By: SCIREWebVideo	https://www.youtube.com/watch?v=XBWfSvq78bE	1:37
3.	5 Guidelines for Addressing Sexual Health after Spinal Cord Injury By: SCIREWebVideo	https://www.youtube.com/watch?v=gx4sryINCQU	3:17
4.	Sexual Health and Spinal Cord Injury: The 3-Step Method By: SCIREWebVideo	https://www.youtube.com/watch?v=VbV39Y3cl7M	2:37
5.	Sexual Health and Spinal Cord Injury: Sexual Self-esteem By: SCIREWebVideo	https://www.youtube.com/watch?v=bF7Lc1K-6Ro	3:31
6.	Spinal Cord Injury: A Sexual Health Framework By: SCIREWebVideo	https://www.youtube.com/watch?v=4XCKzbJDcC	1:18
7.	The PLISSIT model: 4 levels of Intervention for Treating Sexual Health problems By: SCIREWebVideo	https://www.youtube.com/watch?v=vJ6RUWBCuS4	4:15
8.	Sex After Spinal Cord Injury By: FacingDisability	https://youtu.be/QhXLuJNN8t4	1:02

indicator; [Appendix A](#)) following the training. Additionally, barriers to implementation of the self-assessment tool were noted in order to inform the Working Group.

Pilot study methods – outcome indicator

Second, the feasibility of the outcome indicator (sexual health patient questionnaire) was also tested at a tertiary SCI rehabilitation hospital using rapid PDSA cycles. Components from standardized questionnaires (e.g. sexual attitude and information questionnaire, female sexual function index, sexual health inventory for men, etc.) were used to develop the sexual health patient questionnaire. Consecutive patients attending routine follow up appointments were approached by a trained evaluator to complete the questionnaire. Patients were asked if they were comfortable answering sexual health questions prior to consenting to commence with the questionnaire. Physical assistance was provided in completing the paper and pen questionnaire for individuals who could not complete it themselves (i.e. individuals with limited hand functioning). The primary goal of the PDSA cycles for the Sexual Health Domain was to pilot the feasibility of the questionnaire (outcome indicator; [Appendix B](#)). The questionnaire was designed to collect indicator data regarding sexual health patient outcomes (i.e. sexual expression, sexual self-esteem, sexual activity, and sexual satisfaction) and sexual health service delivery (i.e. quantitative data on information provided by SCI/D rehabilitation staff, staff knowledge; and quantitative and qualitative data on sexual health education/information needs). Furthermore, the feasibility and challenges of implementing the questionnaire were noted in order to inform the Working Group.

Results

The selection and refinement of structure, process and outcome indicators related to Sexual Health were primarily driven by the impetus to create a permissive environment where HCPs support and

facilitate individuals with SCI/D to see themselves as sexual beings entitled to sexual satisfaction (as depicted in [Figure 1](#) and the construct definition). Regarding the outcome indicator, standardized questionnaires (e.g. sexual attitude and information questionnaire, female sexual function index, sexual health inventory for men, etc.) were used to develop the sexual health patient questionnaire for the current Project. [Table 2](#) summarizes the denominators, type of indicator and timing of measurement for each of the indicators selected by the Sexual Health Working Group.

Sexual Health indicator piloting

Pilot study results – structure indicator

The pilot data regarding the level of sexual health knowledge includes results from 26 SCI/D rehabilitation HCPs who completed the structure indicator pretest and 19 HCPs who completed the post-test self-assessment tool. There was an increase in the level of knowledge as evaluated by the post-test self-assessment tool following independent review of the video module and workshop attendance. Feasibility issues and challenges experienced included: (1) concerns regarding the confidentiality and privacy of participants' assessment of knowledge; and (2) difficulty completing the tool in electronic form (i.e. majority of the participants completed a printed hard copy of the tool).

Pilot study results – outcome indicator

The pilot data on sexual health patient outcomes and sexual health service delivery includes results from 20 individuals with SCI/D who completed the questionnaire. Questionnaires were completed within an average of 5 min. Two individuals declined participation due to feeling uncomfortable answering personal questions related to their sexual health. The majority of the participants reported that they strongly felt their injury interfered with their sexual activity, sexual interest and sexual self-esteem, with most being unsatisfied with the quality of their sexual life. With respect to the sexual

Table 2 Selected structure, process and outcome indicators for the Sexual Health Domain.

Indicator	Denominator	Type	Time of measurement
Proportion of SCI/D rehabilitation program staff (regulated health care professionals) who have completed the sexual health video module + SCI-High self-assessment tool	Total number of health care professionals within the tertiary SCI rehabilitation team	Structure	Annual
Proportion of SCI/D inpatients having a documented introduction to available local sexual health resources	Total number of SCI/D inpatients per fiscal year	Process	Rehabilitation admission
SCI-High Sexual Health Questionnaire	Total number of SCI/D patients with English proficiency	Outcome	18 months post rehabilitation admission

health service delivery, the majority of the participants reported they were not provided with sexual health information by HCPs and that the level of staff knowledge and/or comfort was not clear. Several participants emphasized the need for timely and tailored information about sexual health (particularly fertility, treatments, counseling therapies and online resources). Feasibility issues and challenges included: (1) the impact of cognitive and motor deficits on the ability to self-administer the questionnaire; (2) inability to understand certain aspects of the questionnaire (i.e. clarifications needed on specific language [e.g. “sexual expression”], and timeframe/setting of sexual health information provided by HCP [e.g. information provided when individual was newly injured or currently, during inpatient or outpatient rehabilitation]); (3) the impact of language barriers (i.e. not proficient in English) on the ability to complete the questionnaire; (4) inability to recall if they were provided with sexual health education/information (i.e. those who were several years post-injury); (5) feeling uncomfortable to discuss personal sexual health information, particularly with an evaluator of the opposite sex; and (6) questionnaire was not applicable considering the current sexual health preferences (e.g. due to age, neurological level of injury, other comorbidities, marital/family status).

Discussion

The onset of SCI/D results in complex biological and psychosocial changes that directly and secondarily negatively impact an individual’s overall quality of sexual life post SCI/D. Although improving sexual sense of self (i.e. perception/expression of oneself as a sexual being) and sexual activity improves quality of life after SCI/D,⁴ sexual health issues are poorly integrated into the current SCI/D rehabilitation care process encompassing both inpatient and outpatient settings.^{4,14–16} The structure, process and outcome indicators selected and developed by the SCI-High Sexual Health Working Group are intended to: (1) identify learning gaps, and improve capacity amongst regulated HCPs in sexual health; (2) articulate SCI/D patients’ sexual health needs; (3) identify service delivery gaps, and better integrate sexual health into the SCI/D rehabilitation care process in order to create a permissive environment that enables open discussion and individual health inquiry; and, (4) improve patient sexual self-esteem, satisfaction and expression.

It is vital to recognize that in order to advance sexual health rehabilitation care, consumer priorities should first be taken into consideration. In 2004, an important survey of 681 individuals living with chronic SCI/D

rated sexuality as the first (in individuals with paraplegia) and the second (in individuals with tetraplegia) of seven identified major priorities dictating quality of life after SCI/D.¹⁷ This was reiterated again in 2012 by a systematic review of 24 health and life priorities for persons with SCI/D that indicated sexual function, along with motor function, bladder and bowel issues, among the top four functional recovery priorities, and that health and relationships followed as important life domains following SCI/D.¹⁸ A current review¹⁹ of a community sample of people living with SCI/D identified sexual health problems as the most common significant problem after the injury (41%), followed by chronic pain (24%). Indeed, it becomes increasingly important for regulated HCPs to dispel the prevailing myth of the injured individual’s inability to and/or lack of interest in participating in sexual activities, and promote sexual health as a rehabilitation priority. It should also be noted that individuals with SCI evolve in their readiness to gain new information related to their sexual health.²⁰ Therefore, it is essential to ensure sexual health information is provided in a timely manner and tailored to the individual’s (and spouse/partner’s) specific needs as they adapt to the new life post-SCI/D. The first step towards this initiative of addressing the needs of those living with SCI/D is creating a permissive environment to enable open discussion and individual health inquiry.

In order to assist the numerous numbers and disciplines of HCPs in the rehabilitation setting with at least initiating the discussion on sexual health concerns with the patients they care for early in their rehabilitation journey, the Sexual Health Working Group felt there should be an initiative to improve capacity (i.e. sexual health knowledge and skills). In fact, HCPs dealing with SCI/D must address sexual health and sexual treatment needs as part of the routine discussion, with the same frequency and comfort as they discuss neurogenic bowel and bladder dysfunction. Furthermore, similar to urologic or bowel care expertise, referrals to experts can be initiated to manage sexual health issues beyond the team’s expertise, but this relies upon the issues first being identified. Patients expect their physicians and other HCPs to bring up the topic of sexuality,^{21–23} and tend not to do so themselves, thus relegating their sexual concerns as “less important” and placing the onus on the patient to initiate and sustain the conversation. Even if patients have the courage to do so, their efforts are often met with the HCPs’ discomfort, and/or feelings of inadequacy, in terms of training to deal with such issues, or their discomfort talking about sex itself. Indeed, staff

and organizational training and education initiatives regarding sexual health may significantly optimize SCI/D rehabilitation²⁴ and help HCPs meet the sexual health needs of individuals with SCI/D. The indicators developed by the Sexual Health Working Group are intended to establish a new minimum threshold of sexual health knowledge among HCPs, identify available local sexual health resources, and determine the unmet sexual health needs of individuals with SCI/D in the first 18 months post inpatient rehabilitation admission. The Sexual Health Working Group's premise was that by creating a permissive environment and articulating resources and client needs, staff may direct their continuing education hours toward developing/enhancing their knowledge and skills (e.g. understanding sexual impairment specific to SCI/D), and guide patients to available local sexual health resources as appropriate (i.e. referring patients to experts for management of sexual health issues).

A number of potential barriers with the implementation of the proposed indicators should be considered. First, it should be recognized that new terminology has been introduced that will require sustained use to increase clarity for scientists and recipients. Second, elements from existing psychometric measurement tools (e.g. sexual attitude and information questionnaire, female sexual function index, sexual health inventory for men, etc.) were used to develop the outcome indicator (sexual health patient questionnaire) for this Project. We only used items from existing standardized tools to identify gaps in rehabilitation care relating to sexual health, therefore not all constructs were addressed as originally conceived and validated. Third, the final outcome measurement tool was modified substantially due to specific feasibility restrictions, such as the 10 min time constraint for indicator implementation. Therefore, the newly developed outcome indicator requires further validation. Fourth, while the sample size (26 HCPs; 20 patients with SCI/D) may generally be low in a quantitative research study, it is deemed sufficient for multiple quality improvement PDSA cycles for quick qualitative evaluations, feedback and refinement. PDSA improvement projects do not require large samples to determine a gap in system performance. In addition, small samples can provide valuable information as well as establish validity in PDSA cycles as it may reduce inadequate infrequent data cycles due to collecting too much data in a given cycle.²⁵ Lastly, feasibility was a primary driver for the initial indicators due to the strong desire to establish communities united in their willingness to discuss sexual health, sexual activity and sexual self esteem.

Preliminary pilot data suggests that implementation of the Sexual Health indicators is feasible in identifying HCPs' level of sexual health knowledge, determining the availability of sexual health resources and articulating the unmet sexual health needs of individuals with SCI/D living in the community. These initial indicators were intentionally designed to be feasible and to facilitate the measurement of advances in sexual health across SCI rehabilitation settings and sectors.

Conclusion

In summary, successful implementation of the selected Sexual Health indicators will concurrently promote a permissive environment and characterize the sexual health needs of individuals with SCI/D during the first 18 months following initial admission to SCI/D rehabilitation. These indicators will identify: (1) the proportion of SCI/D rehabilitation HCPs who have completed preliminary sexual health training and education; (2) the proportion of SCI/D inpatients' who have a documented introduction to available local sexual health resources; and (3) sexual health patient outcomes, and information/educational needs to enhance the quality of sexual life amongst individuals with SCI/D. The implementation of these structure, process and outcome indicators is a first step towards creating a therapeutic environment that enables open discussion and individual health enquiry, thus, leading to the provision of equitable and optimal care related to sexual health after SCI/D.

Disclaimer statements

Contributors None.

Funding This work is embedded in the larger SCI-High Project funded by the Rick Hansen Institute (Grant #G2015-33), Ontario Neurotrauma Foundation (ONF - 2018 RHI HIGH 1057), and Toronto Rehab Foundation. Funding for open access publication fees was provided by Toronto Rehab Foundation and Coloplast A/S.

Declaration of interest Dr. B. Catharine Craven acknowledges support from the Toronto Rehab Foundation as the Toronto Rehabilitation Institute Chair in Spinal Cord Injury Rehabilitation and receipt of consulting fees from The Rick Hansen Institute. Dr. Colleen O'Connell acknowledges support from Cytokinetics, Mallinckroft, Orion, Biogen, Rick Hansen Institute and ALS Canada. Dr. Colleen O'Connell has also been a receipt of consulting fees from Ipsen, MT Pharma, Tidal, and Canopy.

Acknowledgements

The authors would like to acknowledge the time, energy and expertise of Dr. Mark Bayley, Dr. Matheus J Wiest and Heather Flett from Toronto Rehabilitation Institute – University Health Network, and Dr. Sander Hitzig from St. John’s Rehab Research Program– Sunnybrook Research Institute throughout the indicator development process.

ORCID

Gaya Jeyathevan  <http://orcid.org/0000-0001-5525-3214>

Farnoosh Farahani  <http://orcid.org/0000-0002-3937-7708>

Anita Kaiser  <http://orcid.org/0000-0002-3521-7432>

S. Mohammad Alavinia  <http://orcid.org/0000-0002-5503-9362>

B. Catharine Craven  <http://orcid.org/0000-0001-8234-6803>

References

- 1 Hensel DJ, Nance J, Fortenberry JD. The association between sexual health and physical, mental, and social health in adolescent women. *J Adoles Health*. 2016;59(4):416–21.
- 2 Shpancer N. Why do we have sex? We have sex more for connection than for procreation or pleasure. 2012.
- 3 Elliott S. Sexual dysfunction and infertility in individuals with spinal cord disorders. In: Kirshblum S, Lin VW, (ed.) *Spinal cord medicine*. Third ed. New York, NY: Springer Publishing; 2019. p. 411–35.
- 4 Anderson KD, Borisoff JF, Johnson RD, Stiens SA, Elliott SL. The impact of spinal cord injury on sexual function: concerns of the general population. *Spinal Cord*. 2007;45(5):328–37.
- 5 Anderson KD, Borisoff JF, Johnson RD, Stiens SA, Elliott SL. Spinal cord injury influences psychogenic as well as physical components of female sexual ability. *Spinal Cord*. 2007;45(5):349–59.
- 6 Kreuter M, Siosteen A, Biering-Sorensen F. Sexuality and sexual life in women with spinal cord injury: a controlled study. *J Rehabil Med*. 2008;40(1):61–9.
- 7 Miranda EP, Gomes CM, de Bessa J, Jr, Najjar Abdo CH, Suzuki Bellucci CH, de Castro Filho JE, *et al*. Evaluation of sexual dysfunction in men with spinal cord injury using the male sexual quotient. *Arch Phys Med Rehabil*. 2016;97(6):947–52.
- 8 Soler JM, Navaux MA, Previnaire JG. Positive sexuality in men with spinal cord injury. *Spinal Cord*. 2018;56(12):1199–206.

- 9 Craven BC, Alavinia SM, Wiest MJ, Farahani F, Hitzig SL, Flett H, *et al*. Methods for development of structure, process and outcome indicators for prioritized spinal cord injury rehabilitation domains: SCI-high Project. *J Spinal Cord Med*. 2019;42(Suppl 1): S51–S67.
- 10 Alavinia SM, Hitzig SL, Farahani F, Flett H, Bayley M, Craven BC. Prioritization of rehabilitation domains for establishing spinal cord injury high performance indicators using a modification of the Hanlon method: SCI-High Project. *J Spinal Cord Med*. 2019;42(Suppl 1):S43–S50.
- 11 Mainz J, Bartels PD, Laustsen S. The National Indicator Project to monitoring and improving of the medical technical care. *Ugeskr Laeger*. 2003;163:6401–6.
- 12 WHO. Defining sexual health: report of a technical consultation on sexual health. Geneva: World Health Organization; 2006. p. 28–31 January 2002.
- 13 Langley GJ, Moen RD, Nolan KM, Nolan TW, Norman CL, Provost LP. *The improvement guide: a practical approach to enhancing organizational performance*. Second ed. San Francisco, CA: Jossey-Bass; 2009.
- 14 Elliott S, Hocaloski S, Carlson M. A multidisciplinary approach to sexual and fertility rehabilitation: the sexual rehabilitation framework. *Top Spinal Cord Inj Rehabil*. 2017;23(1):49–56.
- 15 Post MW, van Leeuwen CM. Psychosocial issues in spinal cord injury: a review. *Spinal Cord*. 2012;50(5):382–389.
- 16 Simpson LA, Eng JJ, Hsieh JT, Wolfe DL. Spinal cord injury rehabilitation evidence Scire research T. The health and life priorities of individuals with spinal cord injury: a systematic review. *J Neurotrauma*. 2012;29(8):1548–55.
- 17 Anderson KD. Targeting recovery: priorities of the spinal cord-injured population. *J Neurotrauma*. 2004;21(10):1371–83.
- 18 Simpson LA, Eng JJ, Hsieh JT, Wolfe DL. The health and life priorities of individuals with spinal cord injury: a systematic review. *J Neurotrauma*. 2012;29(8):1548–55.
- 19 New PW. Secondary conditions in a community sample of people with spinal cord damage. *J Spinal Cord Med*. 2016;39(6): 665–70.
- 20 Hess MJ, Hough S. Impact of spinal cord injury on sexuality: broad-based clinical practice intervention and practical application. *J Spinal Cord Med*. 2012;35(4):211–18.
- 21 Hinchliff S, Gott M. Seeking medical help for sexual concerns in mid-and later life: a review of the literature. *J Sex Res*. 2011;48(2–3):106–17.
- 22 Mellor RM, Greenfield SM, Dowsell G, Sheppard JP, Quinn T, McManus RJ. Health care professionals’ views on discussing sexual wellbeing with patients who have had a stroke: a qualitative study. *PLoS one*. 2013;8(10):e78802.
- 23 Stead M, Fallowfield L, Brown J, Selby P. Communication about sexual problems and sexual concerns in ovarian cancer: qualitative study. *BMJ*. 2001;323: 836–7.
- 24 Gill KM, Hough S. Sexuality training, education and therapy in the healthcare environment: Taboo, Avoidance, discomfort or Ignorance? *Sex Disabil*. 2007;25:73–6.
- 25 Etchells E, Ho M, Shojania KG. Value of small sample sizes in rapid-cycle quality improvement projects. *BMJ Qual Saf*. 2016;25(3):202–6.

Appendix A SCI-High Sexual Health Self-Assessment Tool to assess level of knowledge of health care professionals involved in spinal cord injury rehabilitation to inform sexual health training and education (structure indicator).



SCI-High Sexual Health – Staff Self-Assessment Tool for the Structure Indicator

Before completing this tool, please watch the following educational videos on YouTube:

(Total of 8 short video modules on sexuality & reproduction. Total time:~20 minutes)

- <https://www.youtube.com/playlist?list=PLi2Dc1h0G7-vGPsd0dG88fJlsgs2SsAM3>
- <https://youtu.be/QhXLuJNN8t4>

After reviewing the above video, this tool should be filled out by all inpatient and outpatient employees (full-time, part-time, and casual) participating in the provision of sexual health services for patients with spinal cord injury or disease (SCI/D) in your organization. We anticipate this will take approximately 10 minutes to complete. The information collected will be used to inform the quality of rehabilitation care for individuals with SCI/D.

1. Date: / / (YY/ MM/DD)
2. Please provide your initials: _____
3. Please identify your parent organization: *(drop down on survey monkey)*

4. Please identify your profession:
 - Kinesiologist
 - Nurse practitioner
 - Occupational Therapist
 - Occupational Therapy Assistant
 - Peer Mentor
 - Physiotherapist
 - Physiotherapy Assistant
 - Psychologist

Physician:

 - Family Physician
 - Psychiatrist
 - Psychiatrist
 - Urologist
 - Neuropsychiatrist
 - Registered Nurse
 - Personal Support Worker
 - Sexologist
 - Social Worker
 - Other: _____



5. Please identify your type of employment:

- Full-time
- Part-time
 - 1 day/week
 - 2 days/week
 - 3 days/week
 - 4 days/week
- Casual

6. Please identify your gender:

- Male
- Female
- Other
- Prefer not to disclose

Work Experience:

7. How many years (or months) of work experience related to SCI/D do you have? _____ Year(s)
_____ Month(s)
8. On average, how many SCI/D patients did you provide sexual health information to within the last 3 months? _____

Training & Resources:

9. Which education and/or certification courses related to sexual health have you participated in? (select any that apply):
- Sexual health education during clinical program
 - Sexual health professional certification
 - Sexual health workshops
 - Online sexual health education modules
 - Seminars and conferences
 - No formal training
 - Informal training:
 - Other: _____



10. Which sexual health resources do you have access to in your clinical program? (check all that apply)

Health Care Providers:

- Family physician
- Nurse
- Nurse educator
- Occupational therapist
- Physiatrist
- Physiotherapist
- Psychologist
- Psychiatrist
- Sexologist
- Social worker
- Urologist

Additional Resources:

- Educational guidebooks and manuals
- Websites and e-learning modules
- Peer mentor

- Don't know
- Other: _____

Self-Assessment:

11. Sexual Health includes one or more of the following (check all that apply):

- | | |
|--|---|
| <input type="radio"/> Sexual functions such as erection, ejaculation, lubrication & orgasm | <input type="radio"/> Fertility & contraception |
| <input type="radio"/> Sexual adjustment | <input type="radio"/> Abstinence |
| <input type="radio"/> Transfers | <input type="radio"/> Relationships |
| <input type="radio"/> Sensory awareness | <input type="radio"/> None of the above |

12. Sexual adjustment includes one or more of the following (check all that apply):

- | | |
|--|--|
| <input type="radio"/> Self-esteem | <input type="radio"/> Contraception |
| <input type="radio"/> Self-image | <input type="radio"/> Managing associated medical conditions |
| <input type="radio"/> Relating to others | <input type="radio"/> None of the above |
| <input type="radio"/> Managing mood / depression | |



13. Having an intimate relationship can assist in the process of recovery following SCI/D:
- True
 - False
 - Not sure
14. After SCI/D, common sexual health problems reported by women may include (check all that apply):
- Changes in menstrual cycles
 - Ovulation
 - Altered genital sensation
 - Problems with thermoregulation
 - Changes in lubrication
 - Weight gain
 - Reduced fertility
 - None of the above
15. Spontaneous/reflexogenic erections are commonly experienced by male patients during the early stages of rehab after SCI/D:
- True
 - False
 - Not sure
16. After SCI/D, common sexual health problems reported by men may include (check all that apply):
- Altered genital sensation
 - Changes in erection quality
 - Avoidance
 - Inability to ejaculate
 - Reduced sperm quality
 - Problems with thermoregulation
 - Weight gain
 - Reduced fertility
 - None of the above
17. After SCI/D, sexual interest and functioning may be affected by the following factors (check all that apply):
- Bowel and bladder continence
 - Presence or absence of a partner
 - Thermoregulation
 - Spasticity
 - Sexual self-esteem
 - Neuropathic pain
 - None of the above
18. Strategies to help individuals maximize their sexual experience after SCI/D are (check all that apply):
- Working to maximize your own potential
 - Being open-minded about other options
 - Learning energy conservation strategies
 - Adapting to new physical changes
 - Trying online dating
 - None of the above
19. In keeping with best practice, the appropriate time for discussion of sexual health and related SCI/D issues is:
- Early post-injury
 - One year post-injury
 - During annual follow-up appointments
 - When the person asks
 - All the above
 - None of the above



20. All rehab healthcare professionals need to be sexual health experts:
- True
 - False
 - Not sure
21. Clinicians can build rapport with patients to address sexual health concerns by (check all that apply):
- Seeking help from other experts outside the rehab team
 - Being non-judgmental
 - Avoiding making assumptions
 - Requesting permission to talk about the individual's sexual health
 - None of the above
22. In the PLISSIT model for addressing sexual health, the acronym stands for:
- Patient, Leisure Inactivity, Sexual Satisfaction, and Information Technology
 - Permission, Limited Information, Specific Suggestions, and Intensive Therapy
 - Patient, Limited Information, Specific Suggestions, and Information Technology
 - Permission, Leisure Inactivity, Sexual Satisfaction and Intensive Therapy
 - None of the above
23. Patient-centred care to address sexual health concerns may include consultation with which of the following (check all that apply):
- Sexual health clinician
 - Psychologist
 - Psychiatrist
 - Fertility specialist
 - Cardiologist
 - Spiritual care
 - Urologist
 - Peer mentor
 - Family physician
 - Occupational therapist
 - Physiotherapist
 - Psychiatrist
 - Nurse
 - Nurse educator
 - Social worker
 - Other: _____
 - None of the above
24. Did you watch the sexual health video module before completing this self-assessment questionnaire?
- Yes
 - No
 - Partially

Appendix B SCI-High Sexual Health patient questionnaire to assess sexual health patient outcomes (i.e. sexual expression, sexual self-esteem, sexual activity, and sexual satisfaction) and sexual health education/information needs (outcome indicator).



SCI-High Sexual Health - Interview Guide Template for the Outcome Indicator Questionnaire (SCI-High Sexual Health Patient Questionnaire)

This template provides the interviewer with a step-by-step guide on conducting a sensitive interview related to the Sexual Health domain.

This gender neutral/fluid questionnaire is intended for all persons, regardless of their sexual orientation or expression.

Prior to the interview, ask the individual with SCI/D the following questions:

Many people with spinal cord injury experience significant changes to their sexuality. May I ask you some questions regarding the changes you've experienced?

If response is "yes", proceed; if response is "no", thank the individual and end the interview.

We are conducting this interview to advance sexual and fertility rehabilitation care. If you feel uncomfortable answering certain questions, you may skip them and go on to the next question(s).

Before we begin, would you like to fill out the questionnaire yourself, or could I continue asking the questions?

If the individual prefers to self-administer (or prefers family member/caregiver to fill out the questionnaire while he/she responds), then provide the questionnaire and a pen.

If the individual prefers the interviewer to continue with question, then proceed with interview.

**Note whether the questionnaire was self-administered or filled out by the interviewer.*

Familiarize yourself with the following terms used in the questionnaire:

Sexual Activity: sexual acts with self or partner(s), which may or may not include penetrative activity

Sexual Enjoyment: the pleasure derived from sexual activity with self or partner(s)

Sexual Expression: the way in which we present and communicate as a sexual being to the world, through feelings and behaviours around sexual acts, sexual orientation and gender identity

Sexual Interest: refers to sexual urge and/or motivation to seek out and/or engage in sexual activity

Sexual Life: the physical, mental, emotional and social aspects of sexual activity, expression and relationships

Sexual Self-Esteem: a positive regard for and confidence in the capacity to experience one's sexuality in a satisfying and enjoyable way

Sexual Self-Satisfaction: the expression of the combination of positive sexual outcomes and relationship intimacy following sexual activity



SCI-High Sexual Health Patient Questionnaire

Name: _____

Date of evaluation: _____

Gender: Male Female Other

Age: years

Year(s) since injury:

Months since injury:

Impairment: Para Tetra

Severity: Complete Incomplete

1. What is your current interest in sexual activity?

Very High High Medium Low Absent Unknown

2. I feel that my spinal cord injury interferes with my sexual enjoyment.

Strongly Agree Agree Neither Agree nor Disagree Disagree Strongly Disagree Unknown

3. I feel my sexual expression is limited by my spinal cord injury.

Strongly Agree Agree Neither Agree nor Disagree Disagree Strongly Disagree Unknown

4. How often do you currently engage in sexual activity either alone or with a partner?

Very Frequently Frequently Occasionally Rarely Never Unknown

5. Are you, most of the time, satisfied with the quality of your sexual life?

Strongly Agree Agree Neither Agree nor Disagree Disagree Strongly Disagree Unknown

6. Was information given by a healthcare professional during the initial 18 months following inpatient rehabilitation admission in the following areas? N/A

<i>Area</i>	<i>Yes</i>	<i>No</i>	<i>N/A</i>	<i>Unknown</i>
Sexual function (sex drive, erection, lubrication, ejaculation, orgasm)				
Fertility & Contraception				
Management of your health affecting sexuality (incontinence, pain, spasm, fatigue, medications)				
Sexual adjustment (mood, self-esteem, confidence, relating to others)				
Physical barriers (i.e. inability to access sex toys or positioning cushions or dating environments)				
Physical barriers (i.e. inability to access sex toys or positioning cushions or dating environments)				



SCI-High Sexual Health Patient Questionnaire

7. Do you feel the health care provider:

<i>Area</i>	<i>Yes</i>	<i>No</i>	<i>N/A</i>	<i>Unknown</i>
Was easy to talk to				
Was knowledgeable				
Had a good attitude when talking with me				
Used the right language				
Directed me to the appropriate resources				

8. Do you think the services provided have assisted with your sexual adjustment?

- Yes
 No
 Not Provided- go to #9
 Unknown
 N/A

9. I would like to learn more about:

<i>Area</i>	<i>Yes</i>	<i>No</i>	<i>N/A</i>	<i>Unknown</i>
Sexual response (sex drive, erection, lubrication, ejaculation, orgasm)				
Treatments (medical and psychological) for sexual dysfunctions				
Sensory awareness (body mapping, vibrators, etc.)				
Fertility & contraception				
Management of other health issues affecting sexuality (incontinence, pain, spasm, fatigue, medications)				
Access to sexual aids (sexual positioning options, cushions, devices)				
Creating a private, safe place to be sexual				
Sexual adjustment (mood, self-esteem, confidence, relating to others)				
Dating and relationships after SCI				
Counselling therapies (i.e. psychologist, cognitive behavior therapy, mindfulness)				
Online resources for sexual and fertility rehabilitation				

10. How can we improve sexual rehab services in the future? _____